

Isolation and Biochemical Characterization of Phosphorus Solubilizing Bacteria (PSB) from Rhizospheric Regions of Apricot (*Prunus armeniaca*) and Peach (*Prunus persica*)

Zaffar Bashir^{1*}, Mohd Yousuf Zargar¹, Fayaz Ahmad Mohiddin², Zahoor Ahmad Baba¹ and Burhan Hamid¹

¹Division of Basic Sciences and Humanities, Sher-e-Kashmir University of Agricultural Sciences and Technology, Shalimar-190025, Srinagar, J & K, India

²Division of Plant Pathology, Sher-e-Kashmir University of Agricultural Sciences and Technology, Shalimar-190025, Srinagar, J & K, India

*Corresponding author: zaffarsahib@gmail.com

Abstract

The present study was aimed to isolate and characterize the selected phosphorus solubilizing bacteria from rhizospheric regions of Apricot and Peach. Apricot and Peach rhizospheric region samples were collected from the different sites of district Anantnag, Jammu and Kashmir, India. Each sample was enriched in Pikovskayas's solid medium (pH 7.5 at 30°C for 5 days). Out of 35 isolates, 7 isolates showed highest zone of clearance on Pikovskayas's medium and were morphologically and biochemically characterized. These 7 isolates belong to genus *Bacillus* (PSB3, PSB9, PSB13, 14), *Pseudomonas* (PSB10, PSB11) and *Micrococcus* (PSB12).

Keywords: Phosphorus solubilizing bacteria, rhizosphere, *Bacillus* and *Pseudomonas*.

Introduction

Various microfloras were reported to solubilize insoluble phosphorus complexes into solution and make possible for plant use (Tripura *et al.*, 2005). Phosphorus plays pivotal in photosynthesis, respiration, energy storage and transfer, cell division, cell enlargement and some other processes in the plants. It is applied to the soil in the form of chemical fertilizers. Its availability to the plant utilization is limited. As inorganic phosphate, it is immobilized rapidly and becomes unavailable to plants (Akhtar *et al.*, 2010). Among the alternative P sources, the most important is locally available rock phosphate (Khan *et al.*, 2009). Majority of the soils throughout the world are P deficient (Muhammad, 2012). The concentration of bioavailable P in soil is very low getting to the level of 1.0 kg⁻¹ (Goldstein, 1994). Plants absorb P as phosphate anions (HPO⁴⁻) or H₂PO⁴⁻) from soil (Rodriguez and Fraga, 1999). Phosphate anions are highly reactive in the soil and their precipitation in soil is pH dependent. In acidic soils, phosphate anions get precipitated with free oxides and hydroxides of iron and aluminium, but in alkaline soils, calcium is the main element involved in P fixation (Igual *et al.*, 2001). Indian soils are characterized by poor and medium status with respect to available P (Baby, 2002; Li *et al.*, 2003; and Ramanathan *et al.*, 2004).

Phosphorus solubilizing microorganisms (PSM) play a vital role in soil P dynamics and availability of phosphate to plants (Richardson, 2001). There are various types of soil microbes which can solubilize this fixed form of P and make available to plants (Illmer *et al.*, 1995.; Kucey *et al.*, 1989.; Richardson, 2001 and Whitelaw *et al.*, 1999). Many soil microorganisms particularly those present in rhizosphere of plants, are able to solubilize fixed form of P to soluble form and makes it available to plants (Dave and Patel, 2003.; Dubey *et al.*, 1997 and Narayanasamy *et al.*, 1981). The principal mechanism of P solubilization is the production of organic acids and phosphatase enzymes which play key role in phosphatase solubilization in the soil (Surange *et al.*, 1995; Duttan and Evans, 1996; Nahas, 1996; Shankar *et al.*, 2013). This research articles describes isolation and characterization of PSB from rhizospheric soils of Apricot and peach.

Material and Methods

It was a laboratory scale study and whole of the work was undertaken in the laboratories of Division of Plant Pathology SKUAST- Kashmir. Different strains of phosphate solubilizing bacteria (PSB) were isolated from Apricot and Peach rhizospheric soils

Sample preparation

In order to isolate PSB, 10 mixed soil samples were collected from Apricot and Peach rhizospheric soils, from various sites in Anantnag, India. The samples were collected at a depth of 10-30 cm .All the tools used for soil sampling were surface sterilized using 70% ethanol and soil samples were placed in sterile bags, transported to laboratory, stored at 4⁰C and processed within a week.

Isolation and screening of efficient PSB strains

The isolation of phosphorus solubilizing bacteria (PSB) from rhizospheric soils of Apricot and Peach was done by following the serial dilution technique. For isolation, soil samples were serially diluted from 10⁻³ to 10⁻⁶ and inoculated on modified Pikovskayas's agar medium which consists of : glucose 10 g, Magnesium sulphate 0.1g, ferrous sulphate trace, Manganese sulphate trace, Tricalcium phosphate as P source, agar 15 g, distilled water 1L, pH was adjusted to 7 before sterilization, followed by pour plate technique and the 48 h incubation at 30⁰C, discrete colonies showing halo zones were picked up, sub cultured in Nutrient agar slants and then preserved (Pikovskayas's, 1948).

Characterization of PSB Strains

Morphological characters

Suspension of each purified culture was prepared and poured on plates having solid media by spread plate method. The inoculated plates were incubated at 25⁰ C till the appearance of colonies. Morphological characters of colonies like size, shape, color and elevation were measured (Goenadi *et al.*, 2000).

Microscopic Characters

The isolated strains were heat fixed called as smear. Crystal violet was flooded for one minute and washed gently by tape water. Then the smears were exposed to Gram's iodine for one minute and washed and drained carefully. 95% alcohol was applied for 30 seconds and washed. Finally the smears were washed and drained with 0.25% safranin for 30 seconds and examined under microscope. It focused on shape and size of bacteria. Pink colored bacteria were named as Gram negative while purple colored were named as Gram positive

Estimation of phosphate solubilization efficiency

For testing the P solubilizing capability of PSB strains, each PSB culture was poured on Pikovskayas's agar plate containing insoluble tricalcium phosphate. The plates having culture were incubated for 4 days at 28°C. Solubilization index was measured by Edi-Premono formula (Edi- Premono *et al.*, 1996).

$$PSI = \frac{\text{Colony Diameter} + \text{Halo Zone Diameter}}{\text{Colony Diameter}}$$

Biochemical tests

The most efficient PSB strains were characterized by biochemical tests (Smibert and Krieg, 1994).

Identification of bacterial strain

Different tests like biochemical, morphological and physiological of the selected phosphate solubilizing bacterial isolates were performed for identification, as per methods defined in Bergey's Manual of Determinative Bacteriology (Holt, *et al.*, 1994). The isolates belong to genus *Bacillus*, *Pseudomonas*, *Micrococcus*. PSB 8, 9, 13, 14 belong to *Bacillus*, PSB 10, 11 to *Pseudomonas*, and PSB12 to *Micrococcus*.

Statistical analysis:

Statistical comparison of different PSB strains for P solubilization efficiency, PSI was undertaken.




Figure 1: Phosphorus Solubilization on PVK Medium.

Results and Discussion

Isolation, purification and characterization of PSB.

Microorganisms were capable of producing a zone of clearance due to the solubilization of inorganic phosphorus (Das, 1989) and were routinely screened in the laboratory by a plate assay method using Pikovskayas's agar medium (pikoviskayas 1948) or tricalcium phosphate medium (Nautiyal, 1999). The primary screening protocol used for the identification of PSB strains usually depends on the use of tricalcium phosphate as a sole source of P in indicator plates. We decided to use tricalcium phosphate as source of P for screening of PSB isolated from rhizospheric soils of Walnut.

Results indicate significant clear halo zone formation around bacterial colonies on Pikovskayas's agar with tricalcium phosphate as P source, which was in agreement with reports of Chung *et al.*, (2005); Barroso and Mahas, (2005). The isolates were also tested for potassium solubilization on specific medium. According to morphological and biochemical characterization, the microorganisms showing zone of clearance belongs to the bacterial genera, and various biochemical tests were performed.

From the various PSB isolates, 7 best strains were selected on the basis of solubilization index for further characterization. The characterization of PSB was done by the following tests.

Gram staining

Gram test was performed to check whether PSB strains were gram positive or negative. In this test, few strains retained pink color and were categorized as gram negative and few retained purple colour and which were gram positive.

P-Solubilization test

For the solubilization test, PSB strain were cultured on Pikovskayas's agar medium and placed in incubator at 28°C for 4 days. The formation of halo zone around the bacterial colony confirmed them as P solubilizer (Seshadari *et al.*, 2000). All the 7 strains are best P solubilizers (**Table 3** and **Figure 1**).

Effect of PSB on pH of the Medium

All the selected 7 PSB strains, showed maximum changes in pH. Change in pH by PSB isolates in broth medium was determined by using pH meter. Maliha *et al* (2004) and Chen *et al* (2005) reported that PSB strains secrete various organic acids which decrease the pH of the mediums resulting in P- solubilization. The drop of pH in PSB broth culture has been reported by several researchers like Rashid *et al.* (2004) and Panhwar *et al.*, (2009), who found PSB very effective in lowering the pH of the broth medium. The morphological, biochemical and solubilization index is given in **Tables 1 to 3**.

Table.1. Morphological characters of phosphorus solubilizing and mineralizing bacteria.

Isolate s PSB	Pigmentation	Margi n	Gram reaction	Slightly raised	Highly raised	Trans-parent	Opaque	Shape	Spore
PSB8	White	Entire	+	-	+	+	-	Long rods	+
PSB9	Whitish	Entire	+	+	-	-	+	Short rods	+
PSB10	Creamy	Entire	-	+	-	-	+	Short rods	-
PSB11	creamy	Entire	-	+	-	-	+	Short rods	-
PSB12	Creamy	Elevat ed	+	-	+	-	+	Minute cocci	-
PSB13	White	Entire	+	-	+	+	-	Short rods	+
PSB14	White	Entire	+	-	+	+	-	Short rods	+

Table. 2. Biochemical characterization of 7 PSB isolates.

Isolates PSB	C	U	O	Dn	M. R	V P	ST	Ca	A. P	G. P	H ₂ S	Ge I	Na Cl	Su	M	G	Ci t	K sol.
PSB8	-	+	-	-	+	-	+	-	+	+	-	-	-	-	+	-	+	+
PSB9	-	+	+	+	-	+	-	-	-	-	-	+	-	+	-	+	+	+
PSB10	+	-	+	-	+	-	-	+	+	+	-	-	-	+	+	+	+	+
PSB11	+	-	+	-	-	+	-	+	-	+	+	-	-	-	+	-	-	+
PSB12	+	+	+	-	+	-	+	+	+	+	+	+	-	+	+	+	+	+
PSB13	+	-	+	-	+	-	+	+	+	-	-	+	-	+	-	+	+	+
PSB14	+	+	+	-	+	-	+	-	+	-	-	-	-	-	+	+	+	+

C = Catalase test, U = Urea Hydrolysis, O = oxidase, Dn= Denitrification test, MR = Methyl red test, V.P = V.P test, ST = Starch Hydrolysis test, Ca = Casein test, A.P = Acid production, G.P = Gas production, H₂S = H₂S production test, Gel = Gelatine hydrolysis, Cit = citrate test, SU = sucrose, M = Mannital, G = glucose, NaCl = 70 % NaCl, K Sol = Potassium solubilization,+ = positive, - = negative.

Table. 3. Zone of clearance and phosphorus solubilization index of seven isolates.

PSB isolates	Diameter of zone of Clearance(cm)	Colony Diameter(cm)	Phosphorus solubilization index(PSI)
PSB8	0.60	0.50	2.20
PSB9	0.40	0.30	2.33
PSB10	0.70	0.40	2.75
PSB11	0.50	0.30	2.67
PSB12	0.60	0.40	2.50
PSB13	0.60	0.30	3.00
PSB14	0.70	0.40	2.75




Figure 1: Graphical representation of solubilization index

Conclusion

Apricot and Peach rhizospheric soil presented a diverse population of PSBs. All the screened isolates PSB8 to PSB14 were more efficient in P solubilization. Therefore these isolates can be used in the production of biofertilizers in order to improve growth of agricultural crops in P deficient soils, constituting an alternative to the application of P fertilizers.

Acknowledgement

Authors are thankful to division of Plant Pathology Sher-e-Kashmir University of Agricultural Sciences and Technology-Shalimar, Srinagar, Kashmir 190025, India for their keen interest, Scholastic guidance, supervision and encouragement to accomplish this research.

References:

- Akhtar, N., Iqbal, A., Qureshi, M. A and Khan K. H. 2010. Effect of phosphate solubilizing bacteria on the phosphorus availability and yield of cotton (*Gossypium hirsutum*). *Journal of Scientific Research*. 40(1): 15-24.
- Baby, U. I.2002. Biofertilizers in tea. *Planters' Chronicle*. 98: 395-396.
- Barroso, C.B. and Nahas, E., 2005. The status of soil Phosphate fractions and the ability of fungi to dissolve hardly Soluble Phosphates. *Appl. Soil Ecol*. 29: 73–83.
- Chen, Z., Ma, S., and Lio, L. 2008. Studies on phosphorus solubilizing activities of a strain of phosphobacteria isolated from chestnut type soil in China. *Bioresource Technology* 99: 6702-6707.
- Chung, H., Park, M., Madhaiyan, M., Seshadri, S., Song, J., Cho, H. and Sa, T. 2005. Isolation and Characterization of Phosphate Solubilizing bacteria from the rhizosphere of crop plants of Korea. *Soil Biol. Biochem*. 37: 1970–1974.
- Das, A. C. 1989. Utilization of insoluble Phosphates by soil fungi. *J. Indian Soc Soil Sci*. 58: 1208–1211
- Dave, A. and H. H. Patel. 2003. Impact of different carbon and nitrogen sources on phosphate solubilization by *Pseudomonas fluorescens*. *Indian Journal of Microbiology*. 43: 33-36.
- Dubey, S. K., Balasundaram, V. R., Pant, L. M., Jayasheela, N., Kawale, B. R. and Mishra, B. 1997. Effect of phosphate dissolving bacteria applied with rock phosphate on nodulation

- and yield of rain fed soybean under different agriculture climatic conditions. *Journal of Indian Society for Soil Science*. 45: 503-505.
- Dutton, V.M., and Evans, C.S. 1996. Oxalate production by fungi: its role in pathogenicity and ecology in the soil environment. *Can. J. Microbiol.* 42: 881-895.
- Edi-Premono M, Moawad A and Vleck PLG. 1996. Effect of phosphate solubilizing *Pseudomonas putida* on the growth of maize and its survival in the rhizosphere. Indonesian *Journal of Crop Science*. 11: 13-23.
- Goenadi, D. H.; Siswanto and Sugiarto Y. 2000. Bio activation of poorly soluble phosphate rock with Phosphorus solubilizing fungus. *Soil Science Society of America Journal*. 64: 927-932.
- Goldstein, A. H. 1994 Involvement of the quinoprotein glucose dehydrogenase in the solubilization of exogenous phosphates by gram-negative bacteria. p. 197–203. In: *Phosphate in Microorganisms: Cellular and Molecular Biology*. (A. Torriani-Gorini, E. Yagil and S. Silver, eds). ASM Press, Washington, D.C. , pp. 223
- Igual, J. M., Valverde, A., Cervantes, E., and Velázquez, E. 2001. Phosphate solubilizing bacteria as inoculants for agriculture: use of updated molecular techniques in their study. *Agronomie*. 21: 561-568.
- Illmer, P., Barbato, A. and Schinner, F. 1995. Solubilization of hardly-soluble $AlPO_4$ with P-solubilizing microorganisms. *Soil Biology and Biochemistry*. 27: 265-270.
- Holt, J. G., Krieg, N. R., Sneath, P. H. A., Staley J. T. and Williams S. T. 1994. In, Bergy's Manual of Determinative Bacteriology, (9th Ed.), Williams and Wilkins Pub, MD, USA.
- Khan, A. A., Jilani, G., Akhtar, M.S., Naqvi, S.S. and Rasheed, M., 2009. Phosphorus solubilizing bacteria: occurrence, mechanisms and their role in crop production. *J. Agric. Biol. Sci.* 1: 48–58.
- Kucey, R. M. N., Jenzen H. H. and Leggett, M. 1989. Microbial mediated increases in plant available phosphorus. *Advances in Agronomy* 42: 199-228.
- Li, S. T., Zhou, J. I., Uang, H. Y., Chen, X. Q. and Du. C. W. 2003. Characteristics of fixation and release of phosphorus in three soils. *Acta Pedologica Sznica*. 40: 908-914.
- Maliha, R. K., Samina, Najma A, Sadia A and Farooq L. 2004. Organic acids production and phosphate Solubilization by phosphate solubilizing microorganisms under in vitro conditions. *Pakistan Journal of Biological Sciences* 7: 187-196.
- Muhammadi K. 2012. Phosphorus solubilizing Bacteria, occurrence, mechanisms and their role in crop production. *Journal of Research and Environment* 2(1): 80-85.
- Nahas, E. 1996. Factors determining rock phosphate solubilization by microorganism isolated from soil. *World J. Microb. Biotechnol.*, 12: 18-23.
- Narayanasamy, G., Ghosh, S. K. and Sarkar, M. C. 1981. Chemical and mineralogical composition of P-rock deposits occurring in India. *Fertilizer News*. 26: 3-9.
- Nautiyal C. S. 1999. .An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. *FEMS Microbiol Lett* . 182: 265–270
- Panhwar, Q. A, Radziah, O., Sariah, M., and Razi, I. M. 2009. Solubilization of phosphate forms by phosphate solubilizing bacteria isolated from aerobic Rice. *International Journal of Agriculture Biology*. 11: 667-673.
- Pikovskaya, R. I. 1948. Mobilization of Phosphorus in soil in connection with vital activity of some microbial species. *Microbiologiya*. 17: 362–370

- Ramanathan, S., Natarajan, K. and Stalin, P. 2004. Effect of foliar nutrition on grain yield of rice fallow black gram. *Madras Agriculture*. 91: 160-163.
- Rashid, M., Khalil, S., Ayub, N., Alam, S. and Latif, F. 2004. Organic acids production and phosphate solubilization by phosphate solubilizing microorganisms (PSM) under in vitro conditions. *Pak. J. Biol. Sci.* 7: 187–196
- Richardson, A. E., 2001. Prospects for using soil microorganisms to improve the acquisition of phosphorus by plants. *Funct. Plant Biol.* 28: 897–906.
- Rodríguez, H., and Fraga, R. 1999. Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnology Advances*. 17: 319-339.
- Seshadri, S., Muthukum, A. R., Lakshminarasimhan, C. and Ignacimuthu, S. 2000. Solubilization of inorganic phosphates by *Azospirillum halopraeferans*. *Current Science*. 79: 565- 567.
- Shankar, T., Sivakumar, T., Asha, G., Sankaralingam S. and Meenakshi Sundaram V. 2013. Effect of PSB on Growth and Development of Chilli and Maize Plants. *World Appl. Sci. J.* 26(5): 610-617.
- Smibert, R. M. and Kreig, N. R. 1994. *Phenotypic characterization. In: Methods for General and Molecular Bacteriology* (Gerhard, P., Murray, R. G., Wood, W.A. and Krieg, N.R., Eds.). American Society of Microbiology, Washington DC, 611-654.
- Surange, S., Wollum, A. G., Kumar, N. and Nautiyal, C. S. 1995. Characterization of *Rhizobium* from root nodules of leguminous trees growing in alkaline soils. *Can. J. Microbiol.* 43: 891-894.
- Whitelaw, M. A., Harden, T. J. and K. R. Helyar. 1999. Phosphate solubilisation in solution culture by the soil fungus *Penicillium radicum*. *Soil Biology and Biochemistry*. 31: 655-656.